



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

Dissolved gas and isotopic tracers of denitrification

M. J. Singleton, J. E. Moran, B. K. Esser, W. W. McNab, S. F. Carle, B. D. Cey

February 29, 2008

Goldshmidt
Vancouver, Canada
July 13, 2008 through July 18, 2008

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.

Dissolved gas and isotopic tracers of denitrification

M.J. SINGLETON, J.E. MORAN, B.K. ESSER, W.W. McNAB, S.F. CARLE, B.D. CEY¹²³

We present results from field studies in California (USA) where tritium-helium age dating is used in conjunction with major gases (N_2 , O_2 , CH_4 , CO_2), noble gases (He, Ne, Ar, Kr, Xe), and stable isotopes ($^{15}N/^{14}N$, $^{18}O/^{16}O$) in order to document nitrate loading and denitrification associated with confined animal agricultural operations and septic systems. Preliminary results show that in-field extraction of the full suite of dissolved gases will be possible using a new Gas Extraction System under development to augment the current Noble Gas Mass Spectrometry and Membrane Inlet Mass Spectrometry techniques.

Ascribing observed groundwater nitrate levels to specific current and past land use practices is often complicated by uncertainty in groundwater age and the degree and locus of denitrification. Groundwater age dating at dairy field sites using the 3H - 3He method indicates that the highest nitrate concentrations (150-260 mg/L- NO_3) occur in waters with apparent ages of <5 yrs, whereas older waters contain excess N_2 from saturated zone denitrification [1]. At a residential septic system site in Livermore, CA, waters with young apparent ages (<1 yr) proximal to leach line drainage have lower nitrate concentrations and elevated nitrate $\delta^{15}N$ and $\delta^{18}O$ values consistent with denitrification, but little evidence for excess N_2 , indicating that denitrification is occurring in the unsaturated zone.

Degassing of groundwater can complicate efforts to calculate travel times [2] and to quantify denitrification. Degassed groundwater underlying dairy operations is formed by two distinct mechanisms: (1) recharge of manure lagoon water affected by biogenic gas ebullition [3] and (2) saturated zone denitrification producing N_2 gas above solubility in groundwater. Gas loss due to both mechanisms is evident in the concentrations of noble gases and major gases in dairy groundwater samples.

This work performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344

[1] Singleton, M. J., Esser, B. K., Moran, J. E., Hudson, G. B., McNab, W. W., and Harter, T., 2007. *ES&T* **41**, 759-765. [2] Visser, A., Broers, H. P., and Bierkens, M. F. P., 2007. *WRR* **42**, W10434 (1-14) [3] McNab, W. W., Singleton, M. J., Moran, J. E., and Esser, B. K., 2007. *ES&T* **41**, 753-758.